

GAUGE FIELDS WITH QUAZINILPOTENT GAUGE GROUP.

Arthur M. Aslanyan
Kazan State University, Russia

Abstract

We investigate non-linear generalization of Maxwell theory of electromagnetic field keeping the gauge invariance of Lagrangian. New theory, which is standard Yang-Mills theory, is based on Harmonic Oscillator $HO(N, R)$ gauge group. It's a solvable Lie group with nilpotent normal subgroup of codimension 1. We write down the Yang-Mills equation and point out their peculiarities and connection with standard Maxwell theory.

Field equations and interpretation.

We shall operate with so-called Harmonic Oscillator Lie algebra $ho(4, R)$, which is quaziniipotent and admits non-degenerate invariant bilinear form:

$b : L \times L \rightarrow R$, such that $\forall x, y, z \in L$ the following condition:

$$b(ad_z(x), y) + b(x, ad_z(y)) = 0$$

is hold true.

Consider a phase space of harmonic oscillator with generalized variables: p, q and Hamiltonian: $H = \frac{1}{2}(p^2 + q^2)$. It's known that a set $\langle 1, p, q \rangle$ forms a Lie algebra in respect to Poisson brackets:

$$\begin{cases} [1, p] = [1, q] = 0, \\ [p, q] = 1, \end{cases}$$

which is a nilpotent Lie algebra called Heisenberg algebra.

If we join them a Hamiltonian the new set $\langle 1, p, q, H \rangle$ will form new algebra:

$$\begin{cases} [1, p] = [1, q] = 0, \\ [p, q] = 1, \\ [H, p] = q, \\ [H, q] = -p, \end{cases}$$

which is a quazinilpotent Lie algebra and contains Heisenberg algebra as an ideal. Indeed it's a semi-simple sum of 3-dimensional Heisenberg algebra $\langle 1, p, q \rangle$ and one-dimensional algebra $\langle H \rangle$. This algebra was generalized in papers [1](See also [2]). The main property of this algebra is that it admitts non-degenerate invariant bilinear form:

$$\langle u, v \rangle = u^s v^s + u^1 v^4 + u^4 v^1 + z u^4 v^4$$

Here $s, r, t, \dots = \overline{(2, 3)}$, z — any real constant and u^1, u^s, u^4 — are the components of the element u of Lie algebra in the basis $\langle 1, p, q, H \rangle$:

$$u = u^1 1 + u^2 p + u^3 q + u^4 H.$$

Due to the existence of this form we can construct for the gauge fields $\hat{A}_\alpha$ a Lagrangian $L = \frac{1}{4} \langle \hat{F}_{\alpha\beta}, \hat{F}^{\alpha\beta} \rangle$ which extremals are the Yang-Mills equations exactly!

Here hat $\hat{A}_\alpha$ means that it belongs to the matrix representation of gauge Lie algebra, and $\alpha, \beta, \gamma, \dots$ — are the indices on the bundle (Minkowski) manifold.

$\hat{F}_{\alpha\beta} = \partial_\alpha \hat{A}_\beta - \partial_\beta \hat{A}_\alpha + [\hat{A}_\alpha, \hat{A}_\beta]$ — curvature tensor of gauge field $\hat{A}_\alpha$,

Yang-Mills equations are:

$$\begin{cases} \partial^\alpha F_{\alpha\beta}^1 + \omega_{st} A^{s\alpha} F_{\alpha\beta}^t = 0, \\ \partial^\alpha F_{\alpha\beta}^s + 2\omega_{st} A^{[t} F_{\alpha\beta}^{s]} = 0, \\ \partial^\alpha F_{\alpha\beta}^4 = 0, \end{cases}$$

coupled with system

$$\begin{cases} F_{\alpha\beta}^1 = 2\partial_{[\alpha} A_{\beta]}^1 + \omega_{st} A_\alpha^s A_\beta^t, \\ F_{\alpha\beta}^s = 2\partial_{[\alpha} A_{\beta]}^s + 2\omega_{st} A_\alpha^4 A_\beta^t, \\ F_{\alpha\beta}^4 = 2\partial_{[\alpha} A_{\beta]}^4. \end{cases}$$

Here $\omega_{st} = -\omega_{ts}$. The system is semi-splitted in three parts. We see that A^4 component is a pure Maxwell field. We substitute this field to the

second part and find the A^s components. The first part can be rewrighted as a Maxwell equations with sources:

$$\begin{cases} \varphi_{\alpha\beta} = 2\partial_{[\alpha}A_{\beta]}^1, \\ \partial^\alpha\varphi_{\alpha\beta} = J_\beta. \end{cases}$$

where $J_\beta = -\omega_{st}\partial^\alpha(A_\alpha^sA_\beta^t) - \omega_{st}A^{s\alpha}F_{\alpha\beta}^t$ — "sources".

Note that center of $ho(N, R)$ is $< 1 >$ and from physical point of view it has no useful information, because the element 1 was coupled to the set p, q to complete it to the Lie algebra, what is simply a mathematical trick. It brings to the idea that component A^1 don't represent the real physical field, and putting it to be trivial we can regard the first equation like a constrain on sources, more precisely on physical fields A^2, A^3 .

To wright down the gauge transformations for this theory we have first to construct a Lie group with $ho(4, R)$ Lie algebra. We call this group $HO(4, R)$ a harmonic oscilator Lie group [4]. There are two types of gauge transformations (connected with semi-simple splitting of algebra into two subalgebras):

First type:

$$\begin{cases} A^1 \rightarrow A^1, \\ A^s \rightarrow (e^{-\lambda^4\omega})_t^s A^t, \\ A^4 \rightarrow A^4 + \partial\lambda^4. \end{cases}$$

Transformation for A^4 is usual gauge transformation for Maxwell field (generally for any abelian gauge field). And transformation for A^s are simply rotation in the p, q plane at angle λ^4 :

$$\begin{cases} A^2 \rightarrow \cos(\lambda^4)A^2 + \sin(\lambda^4)A^3, \\ A^3 \rightarrow -\sin(\lambda^4)A^2 + \cos(\lambda^4)A^3. \end{cases}$$

Second type:

$$\begin{cases} A^1 \rightarrow A^1 + \partial\lambda^1 + \omega_{st}\partial\lambda^s\lambda^t + (\lambda^s\lambda^s)A^4, \\ A^s \rightarrow A^s + \partial\lambda^s + \omega_{st}\lambda^tA^4, \\ A^4 \rightarrow A^4. \end{cases}$$

Conclusion. We see that putting p, q — components of gauge field to zero we come to standard Maxwell theory for A^4 — component. Then there are two aspects of this theory.

The local is connected with the physical meaning of p, q — components which satisfy linear equations and non-linear constraints. These components are not independent because of the gauge transformations and we can well put one of them (say A^3) to zero, providing the gauge for

the usual Maxwell field A^4 being fixed. In this case the constraints will be satisfied automatically and we come to the ordinary two-component Maxwell theory with linear constraints on one of the component.

Another aspect is global. We know that in the Maxwell theory all regular in R^3 monopole solutions are trivial. It turns out that the same is hold true for $HO(4, R)$ -theory, but it's much more difficult to prove this [3]. In Maxwell theory we are forced to regard singular monopole solutions (like Dirac magnetic monopole), but it's energy is infinite. Does it valid for the $HO(4, R)$ – theory? We just can claim the same thing in the case when Maxwell sector is trivial ($A^4 = 0$) [3], but it's not clear in the general case.

And the last question: is it possible for non-trivial A^s, A^4 field to give a zero energy? In this case we can speak about energyless electrodynamics, which can lead to the interesting physical effects. to

References

- [1] Gavrilov S.P., Exact solution of Yang-Mills equations with Generalized Harmonic Oscillator gauge group, In Theory of Relativity and Gravitation, V.27, Publishing of Kazan State University, Kazan, 1986
- [2] Aslanyan A.M., Left-invariant metrics on four-dimensional solvable Lie groups, Proceedings of International Conference Geometrization of Physics, Kazan, 1993
- [3] Aslanyan A.M., Asimptotically flat monopole solution of Bogomolny equations with solvable gauge group, Izv. Vuzov. Mathematics (to be published)
- [4] Aslanyan A.M., Gavrilov S.P., Matrix realization of Generalized Harmonic Oscillator Lie group, Izv.Vuzov.Mathematics (to be published)